Journal of Surgical Oncology

Table of Contents

Volume 88, Number 1, October 1, 2004

Guest Editorial: Isolated Limb Infusion for Melanoma: A Simple but Effective Alternative to Isolated Limb Perfusion
John F. Thompson and Peter C.A. Kam 1

Omitting Axillary Lymph Node Dissection in Sentinel Node Negative Breast Cancer Patients Is Safe: A Long Term Follow-Up Analysis
Hans Torrenga, Hans Fabry, Joost R.M. van der Sijp, Paul J. van Diest, Rik Pijpers, and Sybren Meijer 4
The rate of axillary recurrences is evaluated in sentinel node negative patients without further axillary lymph node dissection.

Commentary
Omgo E. Nieweg 7

Predictors of Long-Term Survival in pN3 Gastric Cancer Patients
Junuk Kim, Jae Ho Cheong, Woo Jin Hyung, Jianguo Shen, Seung Ho Choi, and Sung Hoon Noh 9
Although classified as having a stage IV disease as a group according to the 5th UICC cancer staging criteria, the pN3 gastric cancer patients appear to be a heterogeneous group with significantly different long-term prognoses. The metastatic lymph node ratio may serve as a more valuable tool to distinguish pN3 gastric cancer patients with different prognoses than the absolute number of metastatic lymph nodes.

Docetaxel and Hyperthermia: Factors That Modify Thermal Enhancement
Faheez Mohamed, O. Anthony Stuart, Olivier Glehen, Muneyasu Urano, and Paul H. Sugarbaker 14
Factors that may modify the thermal enhancement of docetaxel were studied to optimize its clinical use with hyperthermia.

Invasive Cutaneous Fungal Infections Requiring Radical Resection in Cancer Patients Undergoing Chemotherapy
Ravi Radhakrishnan, Michele L. Donato, Victor G. Prieto, Steven R. Mays, Issam I. Raad, and Henry M. Kuerer 21
Invasive fungal infections may develop in patients receiving chemotherapy. This study describes the clinical manifestations, diagnosis, medical management, and radical surgical technique necessary for cure when invasive fungal infection is refractory to standard antifungal systemic therapy.

Detection of Occult Liver Metastases in Colorectal Cancer by Measurement of Biliary Carcinoembryonic Antigen Concentration: A Prospective Study
Jean-Jacques Tuech, Patrick Pessaux, Nicolas Regenet, Alain Daver, Gerard Lorimier, Roberto Bergamaschi, and Jean-Pierre Arnaud 27
The aim of this study was to determine the potential value of biliary CEA assay in the early detection of occult LM from CRC.

Quality of Life After Colon Interposition by Necessity for Esophageal Cancer Replacement
Huib A. Cense, Mechteld R.M. Visser, Johanna W. van Sandick, Angela G.E.M. de Boer, Bas Lamme, Huug Obertop, and J. Jan B. van Lanschot 32
The aim of this study was to assess the quality of life in patients at least 6 months after esophageal cancer resection and colon interposition without signs of recurrent disease.

Preoperative Intraperitoneal Chemotherapy for Patients With Serosa-Infiltrating Gastric Cancer
Masahiko Yano, Takushi Yasuda, Yoshiyuki Fujiwara, Shuji Takiguchi, Hiroshi Miyata, and Morito Monden 39
Serosa-infiltrating gastric cancer poses a high risk for peritoneal recurrence. Preoperative intraperitoneal chemotherapy could be a feasible and safe treatment, which may result in downstaging of the disease.

Salvage Surgical Resection After High-Dose Ifosfamide (HDIF) Based Regimens in Advanced Soft Tissue Sarcoma (ASTS): A Potential Positive Selection Bias—A Study of the Spanish Group for Research on Sarcomas (GEIS)
Joan Maurel, José Buesa, Antonio López-Pousa, Xavier García del Muro, María Jesús Quintana, Javier Martín, Antonio Casado, Javier Martínez-Trufero, Ramón de las Peñas, and Carmen Balañá 44
An assessment was made of the impact of different factors on response rate, time to tumor progression, and overall survival time in patients with locally advanced or metastatic ASTS included in 3 protocols with HDIF.

Commentary: Keeping Abreast of Soft-Tissue Sarcomas
Charles Catton 50

How I Do It: Portal Vein Resection: A Modified Technique for Reconstruction After Pancreaticoduodenectomy
Marcel C.C. Machado, Estela R.R. Figueira, Marcel Autran C. Machado, José Jukemura, José E.M. Cunha, Marcos V. Perini, and Telesforo Bacchella 52

Volume 88, Number 2, November 1, 2004

Does Overweight Impact on the Prognosis of Patients with Renal Cell Carcinoma? A Single Center Experience of 683 Patients
Luigi Schips, Katja Lipsky, Richard Zigeuner, Stefano Gidaro, Michael Salfellner, Peter Rehak, Karl Pummer, and Gerhart Hubmer 57
The possible impact of body mass index on the prognosis of patients with renal cell carcinoma was investigated. Overweight patients were shown to have a better prognosis than normal weight patients in univariate analysis, which, however, was disproved by multivariate analysis.

Commentary
J. Ortiz, P. Ramachandra, R. Zaki, D. Reich, and C. Manzarbeitia 61

Detection of Lymph Node Micrometastasis in Gastric Cancer by MUC2 RT-PCR: Usefulness in pT1 Cases
Hiromichi Sonoda, Kazuo Yamamoto, Ryoji Kushima, Hidetoshi Okabe, and Tohru Tani 63
The incidence of lymph node micrometastases detected by MUC2 RT-PCR was significantly higher than that detected by CEA RT-PCR. The distribution of lymph node micrometastases detected by MUC2 RT-PCR according to tumor location may be useful for planning minimally invasive procedures.

Postoperative Pulmonary Complications Are Associated with Worse Short- and Long-Term Outcomes After Extended Esophagectomy
Shoichi Kinugasa, Mitsuo Tachibana, Hiroshi Yoshimura, Shuhei Ueda, Toshiyuki Fujii, Dipok Kumar Dhar, Takeru Nakamoto, and Naofumi Nagasue 71
Elderly patients with a history of heavy smoking and poor pulmonary function should be regarded as a high-risk group of patients for developing pneumonia. Pneumonia is associated with worse short- and long-term outcomes after extended esophagectomy.

Randomised Phase 2 Trial of SIR-Spheres® Plus Fluorouracil/Leucovorin Chemotherapy Versus Fluorouracil/Leucovorin Chemotherapy Alone in Advanced Colorectal Cancer
Guy Van Hazel, Anthony Blackwell, James Anderson, David Price, Paul Moroz, Geoff Bower, Gluseppe Cardaci, and Bruce Gray 78
A small randomised clinical trial demonstrated that the addition of a single administration of SIR-Spheres® to a regimen of FU/LV chemotherapy significantly increased the response rate, time to disease progression, and overall survival in patients with advanced colorectal liver metastases.

High-Mobility Group Protein 1(Y): Metastasis-Associated or Metastasis-Inducing?
Alice Evans, Thomas W.J. Lennard, and Barry R. Davies 86
Evidence for the role of HMG-1(Y) in metastasis is summarized and critically appraised.

Elevated Expression of Cyclooxygenase-2 in Breast Cancer and Ductal Carcinoma In Situ Has No Correlation with Established Prognostic Markers
Gurpreet Singh Ranger, Andrew Jewell, Valerie Thomas, and Kefah Mokbel 100
Elevated expression of COX-2 occurs in invasive ductal carcinoma of the breast and DCIS, but is not associated with currently accepted prognostic factors.

Immunohistological Evaluation of Single Small Hepatocellular Carcinoma with Negative Staining of Monoclonal Antibody Hepatocyte Paraffin 1
Takaaki Sugiki, Masakazu Yamamoto, Atsushi Aruga, Ken Takasaki, and Masayuki Nakano 104
Some cases of small hepatocellular carcinoma show negative staining for monoclonal antibody Hepatocyte Paraffin 1.

Commentary: Tuech et al.: Detection of Occult Liver Metastases in Colorectal Cancer by Measurement of Biliary Carcinoembryonic Antigen Concentration: A Prospective Study
L. Peter Fielding 108

Letter to the Editor: Clinical Significance of Intramural Extension of Rectal Cancer to the Sphincter Muscles
Marco Caricato, Fabio Ausania, and Roberto Coppola 109

Volume 88, Number 3, December 1, 2004

Editorial: Medical Errors: A Challenge to All Physicians
Walley J. Temple 111

Commentary: Commentary on the Professional Career of John S. Spratt, Jr., MD, MPH
Hiram C. Polk, Jr 112

Dedication: August 4, 2004
John A. Spratt 113

A Systems Approach to Error Prevention in Medicine
Thomas Jeffery Wieman and Eric Andrew Wieman 115

Another Look at Medical Error
William C. Deskin and Robert E. Hoye 122

The Psychology of Error in Relation to Medical Practice
John R. Pani and Julia H. Chariker 130

Nature and Prevention of Errors in Anesthesiology
Michael F. Heine and Carol L. Lake 143

Errors in Laparoscopic Surgery
Earl Gaar 153

Errors in Pathology and Laboratory Medicine: Consequences and Prevention
Sandra C. Hollensead, William B. Lockwood, and Ronald J. Elin 161

Medication Errors in Pediatrics—The Octopus Evading Defeat
Janice E. Sullivan and John J. Buchino 182

Role of the Pharmacist in Reducing Medication Errors
Paul D. Mangino 189

Medical Errors: The Perspective of the Insurer
Thomas James 195

Volume 88, Number 4, December 15, 2004

Surgical Treatment for Gastric Carcinoma in the Elderly
Arianna Coniglio, Guido Alberto Massimo Tiberio, Monica Busti, Giacomo Gaverini, Luca Baiocchi, Tullio Piardi, Maurizio Ronconi, and Stefano Maria Giulini 201
In a series of 50 octogenarians affected by gastric cancer, we observed that the rate of resective and curative resections as well as long-term survival (considering the mortality for gastric cancer) were correlated only to the cancer stage and not affected by the ASA risk, type of gastrectomy, associated resections and nodal dissection.

Is the Chronologic Age a Contra-Indication for Surgical Palliation of Unresectable Periampullary Neoplasms?
G. Nuzzo, G. Clemente, F. Greco, R. Ionta, and F. Cadeddu 206

p53 and DPC4 Alterations in the Bile of Patients With Pancreatic Carcinoma
Desiree Pantalone, Elisbetta Pelo, Barbara Minuti, Irene Giotti, Ernesto Mazza, Massimo Falchini, Bruno Neri, Gabriella Nesi, Lucia Roberta Girardi, Raffaele Pulli, Giovanni Credi, and Francesca Torricelli 210
Pancreatic cancer is still predominantly diagnosed in advanced stages, and 85%–90% of patients are not eligible for surgery at diagnosis. This study was focused on the mutations of p53 and DPC4 detectable in the bile of patients with histologically proven pancreatic cancers. P53 and DPC4 mutations are present in PANin stage 3. The present study aim is to investigate their possible role in clinical practice.

Prognostic Significance of the Age Factor in the Thyroid Cancer: Statistical Analysis
Laura Falvo, Antonio Catania, Salvatore Sorrenti, Vito D'Andrea, Alberto Berni, Matteo De Stefano, and Enrico De Antoni 217

Up-To-Date on Parathyroid Carcinoma: Analysis of an Experience of 19 Cases
Maurizio Iacobone, Franco Lumachi, and Gennaro Favia 223
Parathyroid carcinoma is considered a rare neoplasm, but in our experience it accounts for 4.7% of causes of primary hyperparathyroidism. Survival is variable and difficult to predict: 79% of patients died because of the disease, after a median survival of 29 months. Significant correlations were found between survival and the disease free-interval; between recurrences, age and preoperative calcium levels.

Preliminary Report on Surgical Technique in Hepatic Parenchymal Transection for Liver Tumors in the Elderly: A Lesson Learned From Living-Related Liver Transplantation
Salvatore Gruttadauria, Cataldo Doria, Claudio H. Vitale, Davide Cintorino, Carlo Scotti Foglieni, John J. Fung, and Ignazio R. Marino 229
We performed a retrospective analysis of perioperative mortality, length of hospitalization (LOS) and blood transfused during surgery in two patient groups who underwent liver resection. In Group A, we described a new technique to resect hepatic parenchyma, using ultrasonic surgical aspirator in association with monopolar floating ball cautery, while in Group B, the crushing clamp technique was utilized. We divided the same population group in < 65 and > 65 years old trying to identify differences between the two groups. Results enforced the idea that the elderly can receive such surgical treatment without hesitation.

Simultaneous Repair of Abdominal Aortic Aneurysm and Resection of Unexpected, Associated Abdominal Malignancies
Giulio Illuminati, Francesco G. Calio', Antonio D'Urso, Riccardo Lorusso, Gianluca Ceccanei, and Francesco Vietri 234
Unexpected abdominal malignancies can be found at laparotomy for elective repair of an abdominal aortic aneurysm. In most cases, simultaneous treatment of both conditions will yield good results.

Biological, Histological, and Clinical Impact of Preoperative IL-2 Administration in Radically Operable Gastric Cancer Patients
Fabrizio Romano, Giovanni Cesana, Mattia Berselli, Maria Gaia Piacentini, Roberto Caprotti, Giorgio Bovo, and Franco Uggeri 240
Gastric cancer patients have a severe immunodeficiency, enhanced by major surgery, and can be prevented by preoperative IL-2 administration. Furthermore they have a low rate of TILs phenomenon which is a tumor-host reaction linked to prognosis. Sixty-nine preoperative gastric cancer entered the study and were randomized to receive or not receive interleukin-2 at 9 million IU/daily for 3 consecutive days before surgery. IL-2 administration was well tolerated and prevented the decrease of total and T helper lymphocyte at 14th and 50th postoperative days, which conversely occurred in control group. IL-2 group seemed to have lower postoperative complication and higher overall and disease-free survivals and more over IL-2 enhances TILs in a significant manner.

Adjuvant Hepatic Arterial Infusion After Curative Resection of Colorectal Liver Metastases Using Removable Intra-Arterial Catheters With Shape-Memory Alloy
Takeshi Tono, Toshiyuki Kanoh, Tadashi Ohnishi, Masaki Matsushita, Yoshiaki Nakano, Yutaka Kimura, Takashi Iwazawa, Hiroshi Yano, Jun Okamura, and Takushi Monden 248
A new approach of prophylactic HAI chemotherapy by using a removable reservoir-catheter system is a promising strategy after curative resection of colorectal liver metastases.

Hepatic Arterial Catheterization Combining Interventional Radiological and Laparotomic Approaches
Masashi Watanabe, Wataru Takita, Haruhiro Nakazaki, Hiroyuki Tanemura, Hironori Kaneko, and Kazuo Kobayashi 256
A new method for implantation of a microcatheter-port system in the hepatic artery using combined interventional radiological and laparotomic approaches is described. This method may enable operators to avoid complicated selective coiling and may lower the incidence of hepatic artery occlusion in patients receiving long-term hepatic arterial infusion chemotherapy.

Two Colons-Two Cancers: Paradigm Shift and Clinical Implications
Pascal Gervaz, Pascal Bucher, and Philippe Morel 261

Ultrasound (US) Guided Central Venous Catheterization of Internal Jugular Vein on Over 65-Year-Old Patients Versus Blind Technique
Massimo Cajozzo, Gerlando Cocchiara, Geo Greco, Roberto Vaglica, Tommaso Bartolotta, Lina Platia, and Giuseppe Modica 267

Review Article: Haemostatic Alterations in Colorectal Cancer: Perspectives for Future Treatment
Jakob Lykke and Hans Jørgen Nielsen 269

Letter to the Editor: Re: Verwaal VJ, van Tinteren H, Ruth SV, et al.: Toxicity of Cytoreductive Surgery and Hyperthermic Intra-Peritoneal Chemotherapy. J Surg Oncol 2004; 85:61–67
Paul H. Sugarbaker 276

Author Index to Volume 88 279

Subject Index to Volume 88 281